



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

402. Proposed by C. N. SCHMALL, New York City.

If (x, y) be a double point on the curve $u \equiv f(x, y) = 0$, show that (1) the two branches of the curve will cut orthogonally if

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0;$$

and (2), if this point be made the origin, then the equation of the tangents to the branches will be

$$(y'^2 - x'^2) \frac{\partial^2 u}{\partial x^2} + 2x'y' \frac{\partial^2 u}{\partial x \partial y} = 0$$

where (x', y') are the current coördinates of points on the tangents.

SOLUTION BY C. E. DIMICK, New London, Connecticut.

If (x, y) be a double point of the curve $u \equiv f(x, y) = 0$, $\partial u / \partial x = \partial u / \partial y = 0$ and the two values of dy/dx at the double point must satisfy the quadratic

$$\frac{\partial^2 u}{\partial x^2} + 2 \frac{\partial^2 u}{\partial x \partial y} \frac{dy}{dx} + \frac{\partial^2 u}{\partial y^2} \left(\frac{dy}{dx} \right)^2 = 0.$$

(Todhunter's Diff. Calc., pages 319, 320.)

If the two tangents are perpendicular, the product of their slopes is -1 , and since the product of the roots of the quadratic $a + bx + cx^2 = 0$ is a/c we have

$$\frac{\partial^2 u}{\partial x^2} / \frac{\partial^2 u}{\partial y^2} = -1 \quad \text{or} \quad \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0.$$

The equations of the two tangents will be $y - y_1 = l_1(x - x_1)$, and $y - y_1 = l_2(x - x_1)$, where l_1 and l_2 are the roots of the quadratic given above. Transposing and multiplying, the equation of the two tangents taken together will be

$$(y - y_1)^2 - (x - x_1)(y - y_1)(l_1 + l_2) + l_1 l_2 (x - x_1)^2 = 0,$$

which becomes on substituting the values of the sum and product of the roots of the quadratic

$$(y - y_1)^2 + 2(x - x_1)(y - y_1) \frac{\partial^2 u}{\partial x \partial y} / \frac{\partial^2 u}{\partial y^2} + (x - x_1)^2 \frac{\partial^2 u}{\partial x^2} / \frac{\partial^2 u}{\partial y^2} = 0,$$

which, upon transforming to (x_1, y_1) as origin, becomes

$$y^2 \frac{\partial^2 u}{\partial y^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + x^2 \frac{\partial^2 u}{\partial x^2} = 0.$$

But $\partial^2 u / \partial x^2 = -\partial^2 u / \partial y^2$ as the tangents are perpendicular.

Hence the equation reduces to

$$(y^2 - x^2) \frac{\partial^2 u}{\partial y^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} = 0,$$

the minus sign as given in the problem being incorrect.

Also solved by I. A. BARNETT, C. K. ROBBINS, GRACE M. BAREIS, J. A. BULLARD, F. M. MORGAN, and H. L. AGARD.

MECHANICS.**304. Proposed by B. F. FINKEL, Drury College.**

A spherical shell, inner radius r and outer radius R , has within it a perfectly smooth solid sphere of the same material and with radius $r_1 < r$. If the inner surface of the spherical shell is also perfectly smooth, determine the motion, after the time t , of the shell and sphere down a rough inclined plane, inclination α .